

Mt. Pleasant Ophthalmology, P.A.

DISEASE AND SURGERY OF THE EYE

Dr. Grady's Patients

Medications: The hospital may substitute generics for the following eye drops.

Pred Forte 1% - one drop in the operated eye every 4 hours while awake

Ocuflox - one drop in the operated eye every 4 hours while awake

Timoptic - one drop 5 hours after the surgery

1. Continue to wear the protective shield over your eye whenever you are sleeping or do not have your glasses on until otherwise directed.
2. Clean your eye carefully each morning with a clean gauze or cotton ball until your eye is no longer producing mucous.
3. You may pull the shield back to put your drops in and replace the tape carefully.
4. Care should be taken to avoid sleeping with pressure on the operated eye.
5. Strenuous activity should be avoided for the first week following surgery. A good rule to remember is keep your head above your heart.
6. Squat rather than lean over to pick up objects from the floor.
7. You should resume all medications that you were using prior to going to the hospital.
8. You should have no severe pain at home after surgery. If your eye hurts, please do not hesitate to call me.
9. Please remember to bring this sheet, as well as all your eye medications, to the office when you come for your post operative checkups.
10. You may take 2 Tylenol tablets every 4 hours on the day of your surgery if needed for discomfort.

IF YOU HAVE ANY QUESTIONS, PLEASE DO NOT HESITATE TO CALL US

John L. Grady, M.D. * Prat Itharat, M.D.
1705 BEAUCASTE RD, MT. PLEASANT, SC 29464
TEL: (843)884-2015 FAX: (843)881-7068